

Response to Compagno and Persico*

V Hnizdo

National Institute for Occupational Safety and Health, 1095 Willowdale Road, Morgantown, WV 26505, USA

Abstract

Some mathematically incorrect claims of Compagno and Persico in their reply (2002 *J. Phys. A: Math. Gen.* **35** 8965) to my comment on their recent paper on self-dressing and radiation reaction in classical electrodynamics are pointed out.

Compagno and Persico (CP) have replied [1] to my comment [2] on their paper on self-dressing and radiation reaction in classical electrodynamics [3]. CP acknowledge the main point of the comment, namely that the expression for the time-averaged electromagnetic self-force obtained in [4] for the test charge of a Bohr–Rosenfeld field-measurement procedure and rejected in [5] as incorrect can be obtained also using a formula for the self-force which they derived by different means in [3]. In view of this fact, CP now endorse the expression in question as correct. However, some claims in their reply call for my response.

The expression for the time-averaged self-force on a spherical uniform charge q of radius a obtained in [4] reads

$$\bar{F} = \frac{q^2}{T} \int_0^T dt' Q(t') f(t') \quad (1)$$

where

$$f(t') = -\frac{1}{2a^3}(2 - \chi)(2 - 2\chi - \chi^2)\Theta(2 - \chi) \quad \chi = \frac{T - t'}{a}. \quad (2)$$

Here, the speed of light $c = 1$ and $Q(t)$ is the charge's one-dimensional trajectory, which is subject to the conditions that $Q(t) = 0$ for $t \leq 0$ and $|Q(t)| \ll a$, $|dQ(t)/dt| \ll c$ for $0 < t < T$; $\Theta(x)$ is the Heaviside step function. Instead of the simple closed-form expression (2) for the function $f(t')$, CP counter-proposed in [5] the expression (normalized here to conform with (2)):

$$f(t') = -\frac{2}{3V^2} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} [\delta^{(n+1)}(T - t') - \delta^{(n+1)}(-t')] \langle r^{n-1} \rangle \quad V = \frac{4}{3}\pi a^3 \quad (3)$$

where

$$\langle r^{n-1} \rangle = \int_{|\mathbf{r}_1| < a} d\mathbf{r}_1 \int_{|\mathbf{r}_2| < a} d\mathbf{r}_2 |\mathbf{r}_2 - \mathbf{r}_1|^{n-1} = \frac{72V^2(2a)^{n-1}}{(n+5)(n+3)(n+2)}. \quad (4)$$

*This article is written by V Hnizdo in his private capacity. No official support or endorsement by the Centers for Disease Control and Prevention is intended or should be inferred.

(The above closed-form expression for $\langle r^{n-1} \rangle$ was given subsequently in [6].) My first point here has to be unfortunately of a rather trivial nature. The sign of expression (3) is correct, if, as in the standard notation of elementary calculus, $u^{(n)}[v(x)] \equiv d^n u(y)/dy^n|_{y=v(x)}$. Thus, e.g., $\delta^{(n)}(T-t) \equiv d^n \delta(x)/dx^n|_{x=T-t}$. With this notation, there is no sign misprint in the function $f(t')$ in [5] that CP now want to correct—but if $\delta^{(n)}(T-t)$ meant instead $d^n \delta(T-t)/dt^n = (-1)^n d^n \delta(x)/dx^n|_{x=T-t}$, as CP suggest, then not only the sign but also the factor $(-1)^n$ would be there in error. In either case, the function $f(t')$ is now given incorrectly by the expression (3) of [1]—it either has the wrong overall sign, or the factor $(-1)^n$ there must be omitted.

Contrary to an assertion of CP, I have never claimed that the expression (3) for the function $f(t')$ is incorrect. I have rather pointed out in [6] that, in order to obtain the time-averaged self-force (1), this expression was used incorrectly by CP in the requisite integration. Since this is an integration with finite limits involving high-order derivatives of the delta function, it cannot be performed as simplistically as CP have done, obtaining [5]

$$\bar{F}_{\text{CP}} = -\frac{2\rho^2}{3T} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} [Q^{(n+1)}(T) - Q^{(n+1)}(0)] \langle r^{n-1} \rangle \quad \rho = \frac{q}{V}. \quad (5)$$

This is the ‘exact’ expression for the time-averaged self-force $\bar{F}$ that CP still claim in [1] to be correct, despite the fact that it is inconsistent with their newly adopted approval of the expression (2) for the function $f(t')$. This can be shown by evaluating the time-averaging integral (1) for $\bar{F}$ using the expression (2) for $f(t')$ and the Taylor expansion of the trajectory $Q(t)$ about the point $t = T$:

$$\begin{aligned} \bar{F} &= \frac{q^2}{T} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} Q^{(n)}(T) \int_0^T dt' (T-t')^n f(t') \\ &= \frac{q^2}{T} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} Q^{(n)}(T) \left\{ \alpha_n - \left[a^{n-2} \left(\frac{2\kappa^{n+1}}{n+1} - \frac{3\kappa^{n+2}}{n+2} + \frac{\frac{1}{2}\kappa^{n+4}}{n+4} \right) + \alpha_n \right] \Theta(2-\kappa) \right\} \end{aligned} \quad (6)$$

where

$$\alpha_n = \frac{48(2a)^{n-2}n}{(n+1)(n+2)(n+4)} \quad \kappa = \frac{T}{a}. \quad (7)$$

This is the correct expression for the time-averaged self-force in terms of the derivatives $Q^{(n)}(T) \equiv \lim_{t \rightarrow T-} d^n Q(t)/dt^n$. (Note for completeness that the correct expression for $\bar{F}$ in terms of the derivatives $Q^{(n)}(0) \equiv \lim_{t \rightarrow 0+} d^n Q(t)/dt^n$ is given by equation (25) of [6].) As $\alpha_n = \frac{2}{3}n\langle r^{n-2} \rangle/V^2$, using (6) we can write $\bar{F}$ for $T \geq 2a$ as

$$\bar{F} = \frac{q^2}{T} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} Q^{(n)}(T) \alpha_n = -\frac{2\rho^2}{3T} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} Q^{(n+1)}(T) \langle r^{n-1} \rangle \quad T \geq 2a \quad (8)$$

which happens to be equal to the first of the two terms in the expression (5) of CP for $\bar{F}$. This demonstrates that expression (5) is incorrect; for $T \geq 2a$, (5) can be corrected by dropping its second term, but for $T < 2a$, no such simple correction is possible.

Far from being ‘convenient’ for an ‘exact’ evaluation of the time-averaged self-force $\bar{F}$, formula (3) is a purely formal expression that has no practical application in an integration

with finite limits. Its use by CP has led to the erroneous expression (5) for $\bar{F}$; when CP use it in [1] to prove the equivalence of expressions (2) and (3), they revert the Taylor expansions $\sum_n (-1)^n \delta^{(n+1)}(t) r^n / n!$ to $\delta'(t - r)$ before performing the finite-limit integration.

So far, CP have responded to my criticism [4] of their re-analysis [7] of the Bohr–Rosenfeld field-measurement procedure only by making mathematically incorrect claims. All these involved rather simple mathematical points about which there should have been no need of explicating since my paper [6] of 2000 (where I derived expression (2) using elementary calculus of the delta function), if not already since my comment [4] of 1999 (where I used Fourier transform methods). It is regrettable that such points have deflected from the interesting issues of physics relating to the famous Bohr–Rosenfeld analysis.

- [1] Compagno G and Persico F 2002 Reply to Comment on Self-dressing and radiation reaction in classical electrodynamics *J. Phys. A: Math. Gen.* **35** 8965–67
- [2] Hnizdo V 2002 Comment on Self-dressing and radiation reaction in classical electrodynamics *J. Phys. A: Math. Gen.* **35** 8961–63 (*E-print* physics/0205022)
- [3] Compagno G and Persico F 2002 Self-dressing and radiation reaction in classical electrodynamics *J. Phys. A: Math. Gen.* **35** 3629–45
- [4] Hnizdo V 1999 Comment on Limits of the measurability of the local quantum electromagnetic-field amplitude *Phys. Rev. A* **60** 4191–95 (*E-print* quant-ph/0210074)
- [5] Compagno G and Persico F 1999 Reply to Comment on Limits of the measurability of the local quantum electromagnetic-field amplitude *Phys. Rev. A* **60** 4196–97
- [6] Hnizdo V 2000 The electromagnetic self-force on a charged spherical body slowly undergoing a small, temporary displacement from a position of rest *J. Phys. A: Math. Gen.* **33** 4095–103 (*E-print* math-ph/0005014)
- [7] Compagno G and Persico F 1998 Limits on the measurability of the local quantum electromagnetic-field amplitude *Phys. Rev. A* **57** 1595–603